

0.3 mg; For animals weighing 51 to 100 lbs: 0.4 mg; For animals weighing over 100 lbs: 0.5 mg. Dosage may be gradually increased up to a maximum of five times the suggested dosage. Following parenteral use, dosage may be continued by oral administration of tablets.

(2) *Indications for use.* For the treatment of vomiting and/or diarrhea, nausea, acute abdominal visceral spasm, pylorospasm, or hypertrophic gastritis.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16183, Mar. 25, 2014]

#### § 522.82 Aminopropazine.

(a) *Specifications.* Each milliliter of solution contains aminopropazine fumarate equivalent to 25 milligrams (mg) aminopropazine base.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs and cats*—(i) *Amount.* 1 to 2 mg per pound of body weight, repeated every 12 hours as indicated, by intramuscular or intravenous injection.

(ii) *Indications for use.* For reducing excessive smooth muscle contractions, such as occur in urethral spasms associated with urolithiasis.

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Horses*—(i) *Amount.* Administer 0.25 mg per pound of body weight, repeated every 12 hours as indicated, by intramuscular or intravenous injection.

(ii) *Indications for use.* For reducing excessive smooth muscle contractions, such as occur in colic spasms.

(iii) *Limitations.* Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16183, Mar. 25, 2014]

#### § 522.84 Beta-aminopropionitrile.

(a) *Specifications.* The drug is a sterile powder. Each milliliter of constituted solution contains 0.7 milligrams (mg) beta-aminopropionitrile fumarate.

(b) *Sponsor.* See No. 064146 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount.* Administer 7 mg by

intralesional injection every other day for five treatments beginning about 30 days after initial injury.

(2) *Indications for use in horses.* For treatment of tendinitis of the superficial digital flexor tendon (SDFT) in horses where there is sonographic evidence of fiber tearing.

(3) *Limitations.* Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16183, Mar. 25, 2014]

#### § 522.88 Amoxicillin.

(a) *Specifications*—(1) Each vial contains 3 grams (g) of amoxicillin trihydrate. Each milliliter of constituted suspension contains 100 or 250 milligrams (mg) amoxicillin trihydrate for use as in paragraph (d)(1) of this section.

(2) Each vial contains 25 g of amoxicillin trihydrate. Each milliliter of constituted suspension contains 250 mg amoxicillin trihydrate for use as in paragraph (d)(2) of this section.

(b) *Sponsor.* See No. 054771 in § 510.600(c) of this chapter.

(c) *Related tolerance.* See § 556.38 of this chapter.

(d) *Conditions of use*—(1) *Dogs and cats*—(i) *Amount.* Administer 5 mg per pound of body weight daily for up to 5 days by intramuscular or subcutaneous injection.

(ii) *Indications for use*—(A) *Dogs.* For treatment of infections caused by susceptible strains of organisms as follows: Respiratory infections (tonsillitis, tracheobronchitis) due to *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, and *Proteus mirabilis*; genitourinary infections (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal infections (bacterial gastroenteritis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; bacterial dermatitis due to *S. aureus*, *Streptococcus* spp., and *P. mirabilis*; soft tissue infections (abscesses, lacerations, and wounds), due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*.

(B) *Cats.* For treatment of infections caused by susceptible strains of organisms as follows: Upper respiratory infections due to *S. aureus*, *Staphylococcus* spp., *Streptococcus* spp.,